



For research use only

ISO9001

Dra I

Product	Quantity	Cat. No.	Remarks
Dra I	1,000 unit	EBR-1049	10 unit/μl

Recognition site

5'...T T T | A A A...3'
3'...A A A | T T T...5'

Source

Deinococcus radiophilus.

Reaction Conditions

Buffer 4. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/μl in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μg/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Not sensitive to *dam*, *dcm* or CpG methylation.

Buf4	37°C	65°C		
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	75	75	50	100



For research use only

ISO9001

Dra I

Product	Quantity	Cat. No.	Remarks
Dra I	1,000 unit	EBR-1049	10 unit/μl

Recognition site

5'...T T T | A A A...3'
3'...A A A | T T T...5'

Source

Deinococcus radiophilus.

Reaction Conditions

Buffer 4. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/μl in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μg/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Not sensitive to *dam*, *dcm* or CpG methylation.

Buf4	37°C	65°C		
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	75	75	50	100