



For research use only

ISO9001

## eRec Protein (*E.coli*)

Product	Quantity	Cat. No.	Remarks
eRec Protein ( <i>E.coli</i> )	200 $\mu$ g	EBT-3034	5 $\mu$ g / $\mu$ l

### Description

eRecA Protein (*E.coli*) is involved in the genetic recombination reactions, DNA repair and UV-induced mutagenesis. RecA promotes the strand exchange of single-strand DNA fragments with homologous duplex DNA. The reaction has three distinct steps: (1) RecA polymerizes on the single-strand DNA, (2) the nucleoprotein filament binds the duplex DNA and searches for a homologous region, (3) the strands are exchanged. eRecA is purified from *E.coli* strain carrying an overexpressed RecA gene.

### Concentration & Storage Condition

5  $\mu$ g /  $\mu$ l, Store at -20°C.

### Storage Buffer

20 mM Tris-HCl, pH 7.5, 1 mM DTT, 0.1 mM EDTA and 50% glycerol.

### QC Tests

Activity, exo and endonuclease activity test, SDS-PAGE purity, performance tests.



For research use only

ISO9001

## eRec Protein (*E.coli*)

Product	Quantity	Cat. No.	Remarks
eRec Protein ( <i>E.coli</i> )	200 $\mu$ g	EBT-3034	5 $\mu$ g / $\mu$ l

### Description

eRecA Protein (*E.coli*) is involved in the genetic recombination reactions, DNA repair and UV-induced mutagenesis. RecA promotes the strand exchange of single-strand DNA fragments with homologous duplex DNA. The reaction has three distinct steps: (1) RecA polymerizes on the single-strand DNA, (2) the nucleoprotein filament binds the duplex DNA and searches for a homologous region, (3) the strands are exchanged. eRecA is purified from *E.coli* strain carrying an overexpressed RecA gene.

### Concentration & Storage Condition

5  $\mu$ g /  $\mu$ l, Store at -20°C.

### Storage Buffer

20 mM Tris-HCl, pH 7.5, 1 mM DTT, 0.1 mM EDTA and 50% glycerol.

### QC Tests

Activity, exo and endonuclease activity test, SDS-PAGE purity, performance tests.