




For research use only

Apa LI

ISO9001

Product	Quantity	Cat. No.	Remarks
ApaLI	1,000 unit	EBR-1015	10 unit/ μ l

Recognition site**Source***Acetobacter pasteurianus*.**Reaction Conditions**

Buffer 4 + BSA. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Supplement with 100 μ g/ml BSA. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/ μ l in 10 mM Tris-HCl (pH 7.5), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

No.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by overlapping CpG methylation (Cleavage of mammalian genomic DNA is blocked).



For research use only

Apa LI

ISO9001

Product	Quantity	Cat. No.	Remarks
ApaLI	1,000 unit	EBR-1015	10 unit/ μ l

Recognition site**Source***Acetobacter pasteurianus*.**Reaction Conditions**

Buffer 4 + BSA. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Supplement with 100 μ g/ml BSA. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/ μ l in 10 mM Tris-HCl (pH 7.5), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

No.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by overlapping CpG methylation (Cleavage of mammalian genomic DNA is blocked).

Buf4	37°C	65°C	CpG	BW
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	100	100	10	100

Buf4	37°C	65°C	CpG	BW
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	100	100	10	100