

For research use only

Hae III Methylase

Product	Quantity	Cat. No.	Remarks	
Hae III Methylase	1,000 unit	EBR-3006	10 unit/μℓ	

ISO9001

Description

The Hae III methylase recognizes 5'--GGCC--3' sequences and methylates internal cytosine (C^5) residues..



Source

Recombinant gene from Haemophilus aegyptius.

Applications

Blocking restriction endonuclease cleavage restricted by Hae III methylation.

Reaction Conditions

1x Methylase reaction buffer in the presence of 160 μM S-adenosylmethionine (SAM). Incubate at 37 $^{\circ}\mathrm{C}$.

Concentration & Storage Condition

Store at -20 $^{\circ}$ C. 10 unit/µl in 50 mM Tris-HCl, pH 7.5, 50 mM KCl, 1 mM DTT, 10 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol

Unit Definition

One unit is defined as the amount of enzyme required to fully protect against Hae III cleavage of 1 μg of λ DNA in 1 hr at 37 $^{\circ}$ C.

Heat Inactivation Condition

65℃ for 20 min.

QC Tests

Activity, exo and endonuclease activity test, SDS-PAGE purity, performance tests.



For research use only

ISO9001

Hae III Methylase

Product	Quantity	Cat. No.	Remarks	
Hae III Methylase	1,000 unit	EBR-3006	10 unit/⊯	

Description

The Hae III methylase recognizes 5'--GGCC--3' sequences and methylates internal cytosine (C⁵) residues..

Source

Recombinant gene from Haemophilus aegyptius.

Applications

Blocking restriction endonuclease cleavage restricted by Hae III methylation.

Reaction Conditions

1x Methylase reaction buffer in the presence of 160 μM S-adenosylmethionine (SAM). Incubate at 37 $^{\circ}$ C.

Concentration & Storage Condition

Store at -20 $^{\circ}$ C. 10 unit/ μ l in 50 mM Tris-HCl, pH 7.5, 50 mM KCl, 1 mM DTT, 10 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol

Unit Definition

One unit is defined as the amount of enzyme required to fully protect against Hae III cleavage of 1 μg of λ DNA in 1 hr at 37 $^{\circ}$ C.

Heat Inactivation Condition

65 °C for 20 min.

QC Tests

Activity, exo and endonuclease activity test, SDS-PAGE purity, performance tests.