



For research use only

ISO9001

100bp Step Ladder Marker, Ready-to-use

Product	Conc.	Cat. No.	Remarks
100bp Step Ladder Marker, Ready-to-use	500 µl (100 µg)	EBM-1103	Ready-to-use

Description

The 100bp step ladder marker is a mixture of double-stranded DNA fragments from 100 to 3,000 bp in 100 bp step increment. For easy size reference on the gel electrophoresis, the 500 bp, 1,000 bp, 2,000 bp, and 3,000 bp are made to be more brighter than the other bands. The 100bp step ladder marker is supplied in a ready-to-use format. This ladder marker can be stained with ethidium bromide or any other known DNA staining methods.

Storage Buffer

- Marker DNA : 100 µg in 0.5 ml of 10 mM Tris-HCl, pH8.0, 1 mM EDTA, 5% Glycerol, 0.005% Bromophenol Blue, and 0.005% Xylene Cyanol

Recommended Storage Condition

- 20°C for 2 year
- 4°C for 6 months
- Room temperature (20-25°C) for 2 months


Usage Information

- Concentration : 1,000 ng/5 µl (100 µg/0.5 ml)
- Recommended loading : 2-5 µl (100-200 lanes, ready-to-use)
- Range : 100 bp step increase from 100 bp to 3 kbp

Cautions

- Always use the fresh tip to take out marker solution.
(If you do not, trace amount of contaminated DNases from buffer tank may degrade marker DNA rapidly)
- Don't boil the product.
- Use appropriate % of gels for separation of 100 to 3,000 bp sizes
(1 to 3% agarose gel is recommended)
- Confirm that the concentration of DNA staining dye is optimal before use.
(Breakage or suboptimal concentration of ethidium bromide in gel is a main cause of low estimation of marker concentration or your DNA. 5 ng of DNA should be seen in normal condition)
- Loading volume and concentration should be optimized by gel size, well size, and running length.
- Low sized DNA bands can be gradually disappeared as running is progressing.
(This is because some DNA is getting out from gel to buffer during horizontal electrophoresis, not because the DNA concentration is incorrect. This will be the same for your DNA)
- This product can not be used for a quantitative purpose.


100bp Step Ladder Marker



5 µl/1,000 ng/lane ;

1% agarose in 0.5x TBE, stained with ethidium bromide

Migration Patterns in Different % of Agarose Gels



- 1Kbp DNA Mass Marker (EBM-1009)
- 100bp DNA Mass Ladder Marker (EBM-1010)
- 500bp Step Ladder Marker (EBM-1101)
- 200bp Step Ladder Marker (EBM-1102)
- 100bp Step Ladder Marker (EBM-1103)
- 50bp Step Ladder Marker (EBM-1104)

1.5%, 0.5x TBE Gel

100V constant

EtBr staining

Recommended Gel Percentages for Separation of Linear DNA

Agarose Gel, %	Range of Separation, bp	Polyacrylamide Gel, %	Range of Separation, bp
0.5	1,000 - 30,000	3.5	100 - 1,000
0.7	800 - 12,000	5	80 - 500
1	500 - 10,000	8	60 - 400
1.2	400 - 7,000	12	40 - 200
1.4	200 - 4,000	20	5 - 100
2	50 - 2,000		

DNA Size Migration with Sample Loading Dyes

Agarose Concentration, %	Xylene cyanol FF	Bromophenol blue	Orange G
0.7 - 1.7	~4000 bp	~300 bp	~50 bp
2.5 - 3.0	~800 bp	~100 bp	~30 bp

Composition of Gel Electrophoresis Buffers

Buffer	Working Concentration	Stock Concentration (per Liter)
Tris-acetate (TAE)	1x	20 mM Tris-acetate
		1 mM EDTA
Tris-borate (TBE)	0.5x	45 mM Tris-borate
		1 mM EDTA