



For research use only

ISO9001

Bgl I

Product	Quantity	Cat. No.	Remarks
Bgl I	1,000 unit	EBR-1035	10 unit/μl

Recognition site

5' ...GCCN NNN | NGGC... 3'
 3'... CGGN | NNN NCCG... 5'

Source

Bacillus globigii.

Reaction Conditions

Buffer 3. 50 mM Tris-HCl (pH 7.9), 100 mM NaCl, 10 mM MgCl₂, 1 mM DTT.

Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/μl in 20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 μg/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by some combinations of overlapping CpG methylation (Cleavage of mammalian genomic DNA is blocked).

Buf3	37°C	65°C	CpG	
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	50	75	100	50



For research use only

ISO9001

Bgl I

Product	Quantity	Cat. No.	Remarks
Bgl I	1,000 unit	EBR-1035	10 unit/μl

Recognition site

5' ...GCCN NNN | NGGC... 3'
 3'... CGGN | NNN NCCG... 5'

Source

Bacillus globigii.

Reaction Conditions

Buffer 3. 50 mM Tris-HCl (pH 7.9), 100 mM NaCl, 10 mM MgCl₂, 1 mM DTT.

Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/μl in 20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 μg/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by some combinations of overlapping CpG methylation (Cleavage of mammalian genomic DNA is blocked).

Buf3	37°C	65°C	CpG	
BUFFER	REACTION	INACTIVATION	METHYL BLOCK	BLUE/WHITE

Buffer	1	2	3	4
Activity	50	75	100	50